



## RELATIONSHIP OF BODY WEIGHT WITH LINEAR BODY MEASUREMENTS IN GOATS

Hamayun Khan<sup>1</sup>, Fida Muhammad<sup>2</sup>, Riaz Ahmad<sup>4</sup>, Gul Nawaz<sup>2</sup>, Rahimullah<sup>2</sup> and Muhammad Zubair<sup>3</sup>

<sup>1</sup>Department of Animal Health, NWFP Agricultural University Peshawar, Pakistan

<sup>2</sup>Directorate of Livestock Research and Development NWFP Peshawar, Pakistan

<sup>3</sup>Artificial Insemination Section, Directorate of Livestock and Dairy Development NWFP Peshawar, Pakistan

<sup>4</sup>London Business Academy, London, UK

E-mail: [hamayun\\_dvm@hotmail.com](mailto:hamayun_dvm@hotmail.com)

### ABSTRACT

The present research work was conducted at Livestock Research and Development Station Surezai, Peshawar. Data was collected on 86 goats of different age groups. 44 were male and 42 were female. The mean bodyweight of male in four age groups (04-12,13-18,19-24, 24-36 months and above) were observed as  $18.60 \pm 1.81$ ,  $25.25 \pm 2.76$ ,  $29.86 \pm 1.28$  and  $41.47 \pm 1.63$ kg, respectively while that of female in the same age groups were noted as  $14.50 \pm 1.19$ ,  $21.0 \pm 3.47$ ,  $24.00 \pm 1.25$ ,  $33.95 \pm 4.97$ kg, respectively. The body length of male recorded in these age groups were found to be  $59.60 \pm 0.74$ ,  $64.38 \pm 1.39$ ,  $69.42 \pm 0.29$ ,  $78.15 \pm 0.60$ cm while that of female in these age groups were investigated to be  $58.70 \pm 0.84$ ,  $60.14 \pm 0.50$ ,  $62.16 \pm 0.60$ ,  $69.31 \pm 1.85$ cm, respectively. The mean height at withers in male in four age groups were noted to be  $60.30 \pm 1.16$ ,  $68.25 \pm 1.42$ ,  $72.14 \pm 0.67$ ,  $82.11 \pm 1.02$ cm while that of female in these age groups were recorded to be  $56.50 \pm 1.28$ ,  $63.14 \pm 0.55$ ,  $66.50 \pm 1.18$  and  $71.42 \pm 2.31$ cm, respectively. The mean heart girth in male in these age groups were recorded to be  $59.10 \pm 0.86$ ,  $66.0 \pm 3.19$ ,  $70.29 \pm 1.19$ ,  $79.63 \pm 0.96$ cm, respectively. In case of female mean heart girth was noted to be  $57.60 \pm 0.95$ ,  $61.29 \pm 1.27$ ,  $64.00 \pm 0.68$  and  $70.15 \pm 1.80$ cm, respectively. Bodyweight was correlated with body length (0.49, 0.12, 0.70, and 0.78), height at withers (0.75, 0.54, 0.62, and 0.72) and heart girth (0.64, 0.55, 0.53, and 0.71), respectively. During present investigation male were found heavier ( $p < 0.05$ ) and longer ( $p < 0.05$ ) than female in all age groups. Similarly the heart girth ( $p < 0.05$ ) as well as height ( $p < 0.05$ ) at withers were also bigger in male than those of the female.

**Keywords:** goat, body, length, height, withers, heart, girth, weight.

### INTRODUCTION

Goat is a multi functional animal and plays a significant role in the economy and nutrition of landless, small and marginal farmers in Pakistan. Goat rearing is an enterprise which has been practiced by a large section of population in rural areas. Goats can efficiently survive on available shrubs and trees in adverse harsh environment in low fertility lands where no other crop can be grown. They contribute to livestock industry in terms of milk, meat skin and hair. In comparison with other domestic animals, goats are the victims of prejudice and neglect, but they have nevertheless fulfilled a most useful task in supplying a part of humane population with milk, meat, hair, leather and other products [5]. In spite of their importance, goat has received little scientific attention.

Therefore, the present research study was designed to observe body weight measurement relationship in Beetal goats at Livestock Research and Development Station Surezai, Peshawar. The Beetal breed is used for meat and milk production. The males have long twisting horns. The breed is similar to the Jamnapari but is superior to it in that it is more prolific and more easily adaptable to different agro ecological condition. Males are raised especially as sacrificial animals for slaughter on Eid-Ul-Azha. To increase meat yield from this breed require genetic improvement of its live weight. Proper measurement of this trait, which is often hard in villages due to lack of weighing scale, is requisite for achieving this goal. The need for estimation of the trait from simple

and more easily measurable variable such as linear body measurements therefore arises.

Studies regarding the linear body measurements of goat have been carried in other region of the world and their possible use for estimating the animals live weights (Mason, N. D. [9]; Robinet [15]; Ngere *et al.* [13], Moruppa and ngere [10]; Ibiwoye and Oyatogun [6]; Osinowo *et al.* [14]; Islan *et al.* [7]; Singh and Mishra [16]; Slipper *et al.* [17].

The present study was carried out to establish the relationship between live weight and some body measurements in Beetal goat as a step towards employing such in body weight estimation for selection and other purpose. Results obtained in present study would also be useful and helpful to farmers and animal scientists who are involved in small ruminant's research.

### MATERIALS AND METHODS

86 goats, 44 male and 42 female were used for the present study. The animals were divided in four age groups. Group A (04-12 months) consist of 10 male and 10 female, Group b (13-18 months) consist of 08 male and 07 female, Group C (19-24months) consist of 07 male and 06 female, Group D (25-36months and above) consist of 19 male and 19 female. Among these animals 42 were managed in Livestock Research and Development Station Surezai, Peshawar, while rest of animals were selected form the surrounding area of the station for collection of the data. The animals were released daily for grazing at



8.00am and they remained outside until 2.00 pm. Drinking water was provided ad lib. Animals received routine inspection and dipping; drenching and vaccination were done for herd health maintenance.

In all 86 sets of measurements were obtained for the four variables considered. Body weight was taken using weighbridge and the following linear body measurements were made using the tailor's tape measure as previously used for linear body measurements in Nigerian Red Sokoto goats [1].

Body length (BL) was measured as the distance from the external occipital protuberance to the base of the tail. Height-at-withers (HAW) was measured as the distance from the surface of a platform to the withers and Heart girth (HG) represented the circumference of the chest.

Data collected were classified on the basis of sex and age. Four Age groups (04-12, 13-18, 19-24, 25-36 months and above) were used. Means  $\pm$  SE for the body weight and linear body measurements (BL, HAW, and HG) were calculated using Minitab Statistical software programme. The relationship of bodyweight and linear body measurements were estimated by Pearson correlation. This was done separately for the two sexes in cases of significant sex effect.

## RESULTS

Table-1 summarises the average measurements obtained for the four traits studied. All the values obtained

for linear body measurements were higher in the male than in the females. The mean bodyweight of male in four age groups (04-12,13-18,19-24,24-36months and above) were recorded to be 18.60 $\pm$ 1.81, 25.25 $\pm$ 2.76, 29.86 $\pm$ 1.28 and 41.47 $\pm$ 1.63kg, respectively while that of female in these age groups were observed to be 14.50 $\pm$  1.19, 21.0 $\pm$ 3.47, 24.00 $\pm$ 1.25, 33.95 $\pm$ 4.97kg, respectively. The body length of male in these age groups were recorded to be 59.60 $\pm$ 0.74, 64.38 $\pm$ 1.39, 69.42 $\pm$ 0.29, 78.15 $\pm$ 0.60cm while that of female in these age groups were found to be 58.70 $\pm$ 0.84, 60.14 $\pm$ 0.50, 62.16 $\pm$ 0.60, 69.31  $\pm$ 1.85cm, respectively. The mean height at withers in male in four age groups were noted to be 60.30 $\pm$ 1.16, 68.25 $\pm$ 1.42, 72.14 $\pm$ 0.67, 82.11 $\pm$ 1.02cm, respectively while that of female in these age groups were recorded to be 56.50 $\pm$ 1.28, 63.14 $\pm$ 0.55, 66.50 $\pm$ 1.18and 71.42 $\pm$ 2.31cm, respectively. The mean heart girth in male of four age groups were found to be 59.10 $\pm$ 0.86, 66.0 $\pm$ 3.19, 70.29 $\pm$ 1.19, 79.63 $\pm$ 0.96cm, respectively while that of female in the same age groups were investigated to be 57.60 $\pm$ 0.95, 61.29 $\pm$ 1.27, 64.00 $\pm$ 0.68and 70.15 $\pm$ 1.80cm, respectively. During present investigation male were found heavier ( $p < 0.05$ ) and longer ( $p < 0.05$ ) than female in all age groups. Similarly the heart girth ( $p < 0.05$ ) as well as height ( $p < 0.05$ ) at withers were also bigger than those of the female.

**Table-1.** Mean ( $\pm$ SE) for bodyweight and linear body measurement (cm).

Age (Months)	Sex	No. of Observation	Body weight (kg)	Body length	Height at Withers	Heart Girth
04-12	M	10	18.60 $\pm$ 1.81	59.60 $\pm$ 0.74	60.30 $\pm$ 1.16	59.10 $\pm$ 0.86
	F	10	14.50 $\pm$ 1.19	58.70 $\pm$ 0.84	56.50 $\pm$ 1.28	57.60 $\pm$ 0.95
	M&F	20	16.55 $\pm$ 1.15	59.15 $\pm$ 0.55	58.40 $\pm$ 0.94	58.35 $\pm$ 0.65
13-18	M	8	25.25 $\pm$ 2.76	64.38 $\pm$ 1.39	68.25 $\pm$ 1.42	66.0 $\pm$ 3.19
	F	7	21.0 $\pm$ 3.47	60.14 $\pm$ 0.50	63.14 $\pm$ 0.55	61.29 $\pm$ 1.27
	M&F	15	23.27 $\pm$ 2.18	62.40 $\pm$ 0.94	65.87 $\pm$ 1.03	63.80 $\pm$ 1.85
19-24	M	07	29.86 $\pm$ 1.28	69.42 $\pm$ 0.29	72.14 $\pm$ 0.67	70.29 $\pm$ 1.19
	F	06	24.00 $\pm$ 1.25	62.16 $\pm$ 0.60	66.50 $\pm$ 1.18	64.00 $\pm$ 0.68
	M&F	13	27.15 $\pm$ 1.21	66.08 $\pm$ 1.09	69.54 $\pm$ 1.02	67.38 $\pm$ 1.14
25-36 and above	M	19	41.47 $\pm$ 1.63	78.15 $\pm$ 0.60	82.11 $\pm$ 1.02	79.63 $\pm$ 0.96
	F	19	33.95 $\pm$ 4.97	69.31 $\pm$ 1.85	71.42 $\pm$ 2.31	70.15 $\pm$ 1.80
	M&F	38	37.71 $\pm$ 1.16	73.73 $\pm$ 0.81	76.76 $\pm$ 1.05	74.89 $\pm$ 0.93

Values within each age group with different superscripts differ significantly ( $p < 0.05$ ).

Table-2 describes the correlation of bodyweight with other body measurements. Bodyweight was correlated with body length (0.49, 0.12, 0.70, and 0.78), height at withers (0.75, 0.54, 0.62, and 0.72) and heart girth (0.64, 0.55, 0.53, and 0.71). Results of correlation analysis of sex on the variables are presented in Table-3. The higher correlation coefficient for male goats in most of the cases indicate that on the basis of the dimension of various body lengths the body weight could be predicated more accurately in male as compared to their female

counterpart as the age advances ( $r = 0.824$  for male in 25-36 months of age).

**Table-2.** Coefficient of correlation between the variable.

Age (months)	Variable 1	BW	BL	HAW
04-12	BL	0.485		
	HAW	0.746	0.477	
	HG	0.623	0.551	0.638
13-18	BL	0.115		
	HAW	0.541	0.603	
	HG	0.713	-0.098	0.552
19-24	BL	0.701		
	HAW	0.617	0.813	
	HG	0.525	0.740	0.813
25-36 and above	BL	0.779		
	HAW	0.722	0.893	
	HG	0.708	0.857	0.870

BL = Body length; BW = Bodyweight; HG = Heart Girth; HAW = Height at withers

**Table-3.** Coefficient of correlation between bodyweight (kg) and linear body measurements.

Age (months)	Sex	BL	HAW	HG
04-12	M	0.570	0.671	0.516
	F	0.342	0.769	0.725
	M and F	0.485	0.746	0.638
13-18	M	0.034	0.776	0.759
	F	-0.337	0.050	0.845
	M and F	0.115	0.541	0.552
19-24	M	0.599	0.503	0.130
	F	0.702	0.134	-0.579
	M and F	0.701	0.617	0.525
25-36 and above	M	0.824	0.575	0.524
	F	0.742	0.677	0.751
	M and F	0.779	0.722	0.708

BL = Body length; HG = Heart Girth; HAW = Height at withers

## DISCUSSION

The findings regarding linear body measurements in the present study are in line with previous findings obtained in those reported by Moruppa and Ngere [10] for Red Sokoto goats in similar age groups. The results obtained regarding the body weight and body lengths of goat are lower than previous report on Indian Beetal goat. The body weight in adult male Indian Beetal was reported to be in the range of 50-62kg while in the adult female of the same breed was recorded to be in the range of 35-40kg, respectively. The body length in adult male and female Indian Beetal goat was found to be 86cm and 70.5cm, respectively. Similarly the chest girth was reported in adult male and female of the same breed to be 86 and 73.5cm, respectively [2]. These may be due to the better nutritional and environmental condition. The present finding regarding the body weight are in close vicinity with previous report about the body weight in same indigenous breed. The body weight in adult male and female Beetal goat was reported to be 46 and 37kg, respectively [8].

These values are with no respect to affect the sex of goat on the variable. The high and significant correlation coefficient between height at withers and heart

girth and bodyweight at 04 to 18 months of age suggests that either of this variable or their combination would provide a good estimate for predicting live body weight in Beetal goats at an early age. The higher coefficient of correlation ( $r = 0.638$  and  $r = 0.552$ ) obtained between the two variables attests to this. Similar trend was also reported in Red Sokoto goats [10, 14].

Mukherjee *et al.* [11, 12] and Singh *et al.* [18] reported the highest and significant correlation value of bodyweight with chest circumference in brown Bengal does and grey Bengal goats, respectively. At later stages (from 19-24 months and onward) the body length assumes more importance as an indicator of live weight ( $r = 0.701$ ,  $r = 0.779$ ).

In conclusion, since the body measurements had high correlation with the body weight, this may be used as selection criteria. Earlier reports also indicated that selection based upon the body measurements should improve the meat production [3, 4]. However, further research is needed to investigate the relationship between the body weight with linear body measurements in same and other breeds of goats in different region of the country at different age with maximum number of observations.



## REFERENCES

- [1] Hassan and Ciroma. A.1991 Bodyweight measurements relationship in Nigerian Red Sokoto goats. Department of Animal Science, Usmanur Danfodijo University, Sokoto, Nigeria. [www.fao.org/wairdocs/ILRI/X55208/520bid.htm](http://www.fao.org/wairdocs/ILRI/X55208/520bid.htm).
- [2] Banerjee G. C. 2000. Beetal goat. A text book of Animal Husbandry. 8<sup>th</sup> Ed. Oxford and IBH publishing Co. Pvt. Ltd, New Delhi and Calcutta: pp. 938-939.
- [3] Bhattacharya B., Ghosh T. K., Duttagupta, R. and Maitra, D. N. 1984. Estimation of body weight in Black Bengal goats from body measurements. Indian Veterinary Journal. 61: 406-408.
- [4] Bose, S. and Basu, S. B. 1984. Relationship between body weight-measurement and meat production in Beetal goats. Indian Veterinary Journal. 61: 670-673.
- [5] Fisher. 1983. Goat production in the tropics. Commonwealth Agricultural Bureaux. Farnham House, Farnham, Royal Sussex, UK.
- [6] Ibiwoye, T. I. I. and Oyatogun, M. O. O. 1987. Body weight estimation from measured parameters in sheep and goats in the Kainji Lake Basin, Nigeria. 12<sup>th</sup> Annual Conference, Nigerian Society for Animal Production, March 22-26. Ahmadu Bello University, Shika-Zaria, Nigeria.
- [7] Islam, M. R., Saadullah, M., Howlader, A. R. and Huq, M. A. 1991. Estimation of live weight and dressed carcass weight from different body measurements in goats. Indian Journal of Animal Science. 61(4): 460-461.
- [8] Khan, B. B. and G. B. Issani. 1994. Goat breed of Pakistan. Animal Husbandry by National Book Foundation. pp. 78
- [9] Mason, I. L. (n.d.). The classification of West African Livestock: Cattle. CAB (Commonwealth Agricultural Bureaux), Farnham Royal, Bucks, UK. pp. 10.
- [10] Moruppa, S. M. and Ngere, L. O. 1986. Biometric studies on the Bornu White and Red Sokoto (Moradi) goat breeds. Paper presented at the 11<sup>th</sup> Annual Conference of Nigerian Society for Animal Production, Ahmadu Bello University, 23-27 March. Nigerian Society for Animal Production, Ahmadu Bello University, Shika-Zaria, Nigeria.
- [11] Mukherjee, D. K., Singh, S. K. and Mishra, H. R. 1981. Phenotypic correlations of body weight with body measurements in grey Bengal goats. Indian Journal of Animal Science. 51: 682-694.
- [12] Mukherjee, D. K., Singh, C. S. P. Mishra, H. R. and Nath. 1986. Body weight measurement relationships in brown Bengal does. Indian Journal of Veterinary Medicine 10: 1004-1006.
- [13] Ngere, L. O. Adu, I. F. and Mani, I. 1979. Report of small Ruminant Breeding Subcommittee. NAPRI Bulletin 1. Ahmadu Bello University, Shika-Zaria, Nigeria.
- [14] Osinowo, O. A., Olorunju, S. A. S. Otchere, E. O. and Arigi, L. A. 1989. Development of a weigh-band for Yankasa sheep and Red Sokoto goats. Paper presented at the 14<sup>th</sup> Annual Conference of the Nigerian Society for Animal Production held at Makurdi. 2-6 April.
- [15] Robinet, A. H. 1967. La chèvre Rousse de Maradi son exploitation et sa place dans l'économie et l'élevage de la République du Niger. Revue d'Élevage et de Médecine Vétérinaire des Pays Tropicaux 20(1): 129-186; Animal Breeding Abstracts 35, No. 3746.
- [16] Singh, P. N. and Mishra, A. K. 2004. Prediction of body weight using conformation traits in Barbari goats. Indian Journal of small Ruminants. 10 (2): 173.
- [17] Slippers, S. C., Letty, B. A. and de Villiers, J. F. 2000. Prediction of the body weight of Nguni goats. South African Journal of Animal Sciences. 30 (1): 127-128.
- [18] Singh, N. R. Mohanty, S. C. and Mishra, M. 1987. Prediction of body weight from body measurements in black Bengal goats: a note. Indian Journal of Animal Production and Management. 3: 46-49.