



PERFORMANCE AND ECONOMIC ANALYSIS OF BROILER STARTER CHICKS FED ENZYME SUPPLEMENTED SHEEP MANURE-BASED DIETS

Onu P. N.¹, Madubuike F. N.², Onu D.O.³ and Ekenyem B. U.²

¹Department of Animal Production and Fisheries Management, Ebonyi State University, Abakaliki, Ebonyi State, Nigeria

²Department of Animal Science, Imo State University, Owerri, Imo State, Nigeria

³Department of Agricultural Economics, Management and Extension, Ebonyi State University, Abakaliki, Ebonyi State, Nigeria

E-Mail: nnenwamazi@yahoo.com

ABSTRACT

A 35 day feeding trial was conducted to evaluate the effects of exogenous enzymes supplementation of heat-treated sheep manure-based diets on the performance of broiler starter chicks. 210 7-day old broiler chicks were randomly allotted to seven groups in a completely randomized design. Each group was replicated three times. Seven experimental broiler starter diets were formulated such that diet 1 contained 0% heat-treated sheep manure (HSM). Diets 2, 4 and 6 contained 5%, 10% and 15% HSM without supplementation respectively, while diets 3, 5 and 7 contained 5%, 10% and 15% HSM supplemented with 100mg of exogenous enzyme (Roxzyme G.), respectively. Results show significant ($P < 0.05$) differences in the feed intake, body weight gain, feed conversion ratio, daily protein intake and protein efficiency ratio of the birds. The feed intake of the birds fed 10% and 15% unsupplemented HSM diets were significantly ($P < 0.05$) higher than the other groups. Body weight gain, feed conversion and protein efficiency ratios of birds fed 15% unsupplemented diets were significantly ($P < 0.05$) depressed. Enzymes supplementation of HSM diets significantly ($P < 0.05$) improved the performance of the birds, reduced the cost of feed intake and the cost of feed per kg weight gain over the control.

Keywords: broiler starters, enzymes supplementation, heat-treated sheep manure, performance.

INTRODUCTION

In Nigeria, poultry enterprise is prominent among livestock production and contributes substantial amount of meat and eggs for human consumption. The industry has undergone remarkable growth over the years. For instance, the Nigeria poultry production expanded with the domestic meat production increasing from 172,000 MT in 1999 to 211,000 MT in 2005 (Alabi and Aruna, 2005). However, the expansion is not keeping pace with domestic consumption requirements and is being hampered by fluctuations in the supply of good quality feed.

In Nigeria, poultry feed is based primarily on corn and soyabean meal that constitute the most expensive components of the diets. The scarcity and high cost of these feed ingredients are significant limiting factors to the sustainability and development of poultry production due to the inadequate financial resources available to the farmers. This has led to increasing cost of poultry production, since 70 - 80% of production cost is attributed to feed (Yashim *et al.*, 2007). This limitation imposed by scarcity of feed ingredients has forced animal nutritionists to exploit cheaper and locally available agricultural and agro-by product that can meet the nutrient requirement of poultry (Onu, *et al.*, 2006; Onu, 2007).

One of such agricultural by-products is sheep manure which has always been discarded as waste and unusable by man and any conventional industry. Recently, sheep manure was introduced into monogastric diets by nutritionists as preliminary trials (Abeke *et al.*, 2008; Onu and Otuma, 2008). However, results of the preliminary studies showed that the nutritive value of sheep manure is negatively affected by its low energy and

high fibre contents. Pioneering studies (Abeke *et al.*, 2003; Onu, 2007; Abeke *et al.*, 2008) show that, its high inclusion in poultry diets results in depressed performance of the animals. The present study believes that the sub-optimal performance of the animals fed sheep manure based diets could be overcome by inclusion of exogenous enzymes such as Roxzyme in the diets. Exogenous enzymes have been shown to increase the digestibility of fibrous feed ingredients by disrupting the plant cell walls, and by enhancing nutrients absorption (Acamovic, 2001).

However, the efficiency of exogenous enzymes in diets can only be assessed by evaluating the performance of birds that are fed the diets containing the enzymes compared to those that are not. It was based on this premise that the present study was conducted to evaluate performance and economic analysis of broiler starter chicks fed enzyme supplemented sheep manure based diets.

MATERIALS AND METHODS

The experiment was conducted at the Teaching and Research Farm (Poultry Unit) of the Department of Animal Science, Ebonyi State University, Abakaliki, Nigeria with the approval of the Committee for Animal Experiments of the Institution.

Sources and processing of sheep manure

The sheep manure used in this experiment was sourced from local sheep farmers in Iboko, Izzi local Government Area of Ebonyi State, Nigeria. The manure was oven-dried at 80°C for 3 hours and ground in a hammer mill into a meal suitable for incorporation into



poultry feed. Proximate analysis of the heat-treated sheep manure (HSM) was conducted using standard methods (AOAC, 1995) to determine the percentage crude protein, crude fibre, ether extract and total ash (Table-1).

Table-1. Chemical composition of heat-treated sheep manure.

Nutrients	0%
Dry matter	91.80
Crude protein	21.18
Ether extract	4.95
Crude fibre	21.90
Total ash	32.80
ME (Kcal/kg)	1817.16
Calcium	1.34
Phosphorus	0.56

Experimental diets

Seven experimental starter diets were formulated such that diet T₁ which served as the control contained 0% Heat-treated Sheep Manure (HSM) (Maize - based), while diets 2, 4, and 6 contained 5%, 10% and 15% HSM without enzyme supplementation. Diets 3, 5 and 7 contained 5%, 10% and 15% HSM supplemented with 100mg Roxazyme per kilogram weight of feed respectively. The ingredients were measured out and mixed with a spade on a concrete floor. Turning was vigorously done to ensure good mixing of ingredients and homogeneity.

Experimental animals and procedure

A total of two hundred and twenty (220) day old Anak broiler chicks procured from a commercial hatchery were used for the experiment. Prior to the commencement of the experiment, the birds were fed normal broiler starter chick diets (sheep manure and enzyme - free diet) for 7 days (adaptation period). On the 8th day, two hundred and ten (210) chicks were selected and divided into seven treatment groups of thirty (30) birds each. Each group was randomly assigned to an experimental diet in a completely randomized design. Each group was further sub-divided into three replicates of 10 birds and kept in a compartment measuring 4m x 4m. Heat was provided to the birds using electric bulbs. The chicks were vaccinated against New Castle disease in the first and third weeks and against Gumboro disease in the second and fourth weeks. Prophylactic treatment against coccidiosis with Embazin forte at two weeks of age was also given to the birds.

Fresh water and corresponding diet were provided *ad libitum* throughout the experimental period. Before the commencement of the experiment, the birds were weighed to obtain their initial body weights. The birds were subjected to standard broiler management procedure (FASS.1999). The feeding trial lasted 35 days.

Economics of production

The market cost of the ingredients at the time of the experiment were used to calculate the cost of feed per

kilogram (₦), cost of feed per 100 kilogram (₦), total cost of feed consumed (₦), cost of feed per kilogram weight gain (₦) and cost savings (%).

Statistical analysis

Data obtained were subjected to a one-way analysis of variance while differing treatment means were separated for significance using Duncan's New Multiple Range Test as outlined by Obi (2002).

RESULTS AND DISCUSSION

Chemical composition

Table-1 shows the proximate composition of heat-treated sheep manure and the calculated chemical composition of the experimental broiler starter diets is shown in Table-2.

Heat-treated sheep manure (HSM) contained 91.80% dry matter, 21.18% crude protein, 4.95% ether extract, 21.90% crude fiber, 32.80 ash and 1817 ME (kcal/kg) (Table-6). The use of a feed ingredient in feeding livestock presupposes that the nutritive value in terms of nutrient content and availability be known (Alalade *et al.*, 2006). Fibre is often used as a negative index for measuring the nutritive value of feed ingredients and in the prediction of its total digestible nutrients and net energy (Van Soest *et al.*, 1991). The high crude Fibre content of HSM suggests a low nutritive value of HSM for broilers when included at a high level. However, the dry matter, crude protein, ether extract, ash and ME (kcal/kg) values are slightly higher than the values reported in the works of Abeke *et al.* (2003). The observed variation in the proximate composition is not surprising, as this may have resulted from the chemical composition of the type of pasture consumed by the animals and the processing methods.

The dietary inclusion of HSM at up to 15% in the diet did not alter appreciably the crude protein and ether extract levels of the diets. These levels therefore, met the requirements of the young broiler chicks (NRC, 1994; Aduku, 1993). Increasing HSM inclusion in the diets resulted in a corresponding increase in the crude fibre level of the diets, brought about by the high fibre content of the HSM. It was however observed that the increasing fibre content of the diets resulted in a gradual decrease in the metabolizable energy content of the diets. The ash content of the diets increased progressively as the level of HSM increased.

Performance of broiler chicks

Table-3 shows the performance of the young broiler chicks fed the experimental diets. Feed intake, body weight gain, feed conversion ratio, daily protein intake and protein efficiency ratio of the broiler chickens were affected by the treatments.

The feed intake of the birds fed 10% and 15% unsupplemented HSM diets were significantly ($P < 0.05$) higher than the group on 0% HSM diets. There was no significant variation ($P > 0.05$) among the chicks fed



enzyme supplemented diets and the control in their feed consumption.

The increased feed intake of the birds fed 10% and 15% unsupplemented HSM diets is understandable. The observed effects are attributed to the low energy and high crude fibre levels of the diets. Heat-treated sheep manure is low in energy and high in crude fibre and as the inclusion level increased, the fibre level of the diets also increased. Since the diets are bulky with low energy levels, the birds apparently increased their intake in an attempt to satisfy their energy requirement due to calorie dilution of diets by the 10% and 15% HSM inclusions. This is so because birds would normally adjust their feed intake to meet their dietary energy requirements. This is in agreement with the reports of Onu (2006); Onu *et al.* (2006); Tufarelli *et al.* (2007) and Teimouri *et al.* (2005), who observed that feed intake of birds' increases as the fibre content of the feed increases.

The comparable intake of enzyme supplemented and the control diets may have emanated from the breakdown of the non-starch polysaccharides in the diets by the enzyme, leading to enhanced metabolizable energy value for the enzyme supplemented diets. Oldalle and Hoffman (1996) in an earlier report observed that diets supplemented with enzymes had their metabolizable energy increased by 10%. This finding is also supported by Samarasinghe *et al.* (2000) and Richter *et al.* (1995). They found that feed intake decreased by addition of exogenous enzymes due to the fact that birds fulfill their requirements by taking less amount of feed.

Body weight gain of birds fed 15% unsupplemented HSM diet were significantly ($P < 0.05$) depressed while birds fed 5% enzymatic diet had the highest ($P < 0.05$) body weight gain. The body weight gain of the birds fed the other treatment diets compared favourably ($P > 0.05$) with the control. At the various levels of HSM dietary inclusion, enzyme supplementation of the diets significantly ($P < 0.05$) improved the body weight gains of the birds.

The marked reduction in weight gain at 15% HSM inclusion level without enzyme supplementation despite the higher feed intake of the birds fed this diet indicated that the feed was not adequately translated into increased body weight gain. This is attributed in part to decreased digestibility and utilization of the diet emanating from the high fibre content. Ajaja *et al.* (2003) had observed that high fibre diets decrease nutrient utilization and precipitate metabolic dysfunctions with the attendant weight reduction in monogastric animals and this may have been responsible for the observed weight loss in this study. This observation could be due to the fact that birds mobilized body nutrients for maintenance of metabolism in the place of high dietary fibre and low energy. An (1994) had earlier reported that when birds are starved or when energy levels drop below body requirements, birds tend to mobilize body energy reserves for maintenance, and this may have applied to the observed results in this study.

The enhanced weight gain of the birds on enzymatic diets compared to the birds fed unsupplemented diets suggests that exogenous enzyme supplementation could improve growth in chickens. Response to enzymes could be due to hydrolysis of non-starch polysaccharides and consequent elimination of the negative effects of those polysaccharides on chicks. This may have resulted in greater digestion and absorption of nutrients within the small intestine. Similar observations have been reported by Saki *et al.* (2005); Kongbuntad *et al.* (2006); Onu (2006); Zou *et al.* (2006); Ani and Omege (2007) and Zakaria *et al.* (2008). According to Broz *et al.* (1994) and Zobell *et al.* (2000) exogenous enzymes complements the digestive enzymes of poultry by hydrolyzing the NSPs in cereals, agro-by products and vegetable proteins, thereby decreasing gut viscosity and thus improve nutrient absorption. Feed enzymes also have the ability to alter the bacterial population by digesting the bacteria that colonize the tract and increase the quantity of protein amino acid digested in the pre cecal section of the tract (Bedford, 1997; Gunal and Yasar, 2004) enhancing weight gain of the birds as in this study.

Birds fed 15% unsupplemented (T_6) diet recorded the poorest ($P < 0.05$) feed conversion and protein efficiency ratios. There was marked improvement in feed conversion ratio of birds on supplemented diets over those fed unsupplemented HSM diets at the various levels. The daily protein intake of the birds on diet containing 15% unsupplemented HSM dietary level was significantly higher than those on the other treatment diets. However, this did not differ significantly ($P > 0.05$) from those fed 10% unsupplemented diet. Enzyme supplementation of the diet at different level of inclusion improved the protein efficiency ratio of the birds.

The reduction in feed conversion and protein efficiency ratios of the birds fed 10% and 15% unsupplemented HSM diets has been reported earlier (Abeke *et al.*, 2003). The observed reduction in FCR and PER could be attributed to the deleterious effect of high fibre, which limits the amount of energy and protein available to the birds, and correspondingly contribute to excessive nutrient excretion. However, this limitation was overcome by enzyme supplementation as evident in the superior FCR and PER of the birds fed enzyme treated diets in this experiment. The improved feed conversion ratio achieved with exogenous enzyme supplementation is in agreement with the findings of Jackson *et al.* (2004) and Zou *et al.* (2006) who reported that exogenous enzymes greatly improved the feed conversion ratio of broiler chicks fed enzyme supplemented diets.

Dietary inclusion of heat-treated sheep manure steadily reduced the cost of feed per kilogram as shown on Table-4. There were significant ($P < 0.05$) variations in the cost of feed intake and the cost of feed per kg weight gain of the birds. Enzyme supplementation significantly reduced the cost of feed intake and the cost of feed per kg weight gain over the control. Although 15% HSM inclusion was the least expensive in terms of feed cost per kilogram, feed cost per kilogram weight gain was highest



for birds on 15% unsupplemented HSM diets. This is because of the need to eat more by the broilers to gain 1 kg weight. Thus, the high feed intake, poor feed conversion efficiency and utilization and poor growth rate of the chicks that consumed the 15% unsupplemented HSM diet may have contributed greatly to the high cost of production. Nevertheless, enzyme supplementation significantly reduced feed cost per kilogram weight gain and consequently improved cost saving. The observed

reduction in feed cost / kg weight gain resulting from enzyme supplementation which enhanced cost saving on the production of the birds may probably be due to reduction in feed intake, improved feed conversion efficiency and utilization that resulted to the weight gains of the broilers. Similar results have been reported by earlier studies by Ajaja *et al.* (2003); Onu *et al.* (2006) and Ani and Omeje (2007).

Table-2. Ingredient and chemical composition of experimental broiler starter diets (%).

	T ₁	T ₂	T ₃	T ₄	T ₅	T ₆	T ₇
Ingredients	0.0%	5.0%	5.0%+E	10.0%	10.0%+E	15.0%	15.0%+E
Maize	50.00	47.00	47.00	44.0	44.00	41.00	41.00
SBM	25.00	23.00	23.00	21.00	21.00	19.00	19.00
HSM	0.00	5.00	5.00	10.00	10.00	15.00	15.00
Wheat offal	10.00	10.00	10.00	10.00	10.00	10.00	10.00
Fish meal	5.00	5.00	5.00	5.00	5.00	5.00	5.00
PKC	6.00	6.00	6.00	6.00	6.00	6.00	6.00
Bone meal	3.00	3.00	3.00	3.00	3.00	3.00	3.00
Salt	0.25	0.25	0.25	0.25	0.25	0.25	0.25
Premix*	0.25	0.25	0.25	0.25	0.25	0.25	0.25
Lysine	0.25	0.25	0.25	0.25	0.25	0.25	0.25
Methionine	0.25	0.25	0.25	0.25	0.25	0.25	0.25
Chemical composition							
Crude protein	21.72	21.22	21.22	21.12	21.12	21.04	21.04
Ether extract	4.28	4.34	4.34	4.40	4.40	4.45	4.45
Crude fibre	4.42	5.31	5.31	6.19	6.19	7.08	7.08
Total ash	8.03	9.53	9.53	10.58	10.58	11.68	11.68
ME(Kcal/kg)	2907.90	2839.56	2839.56	2771.22	2771.22	2702.84	2702.84

*To provide the following per kg of feed: Vit. A, 10, 000 IU, vit B₁, 075g, Biotin, 0.05g, Folic acid 1g, Chlorine chloride 250g Copper, 8g, Manganese, 64g, Iron 32g, Zn 40g Iodine 0.6g, Flavomycin 100g, Spiramycin 5g, 3-nitre 50g, DL-methionone, 50g, Selenium, 0.6g, Lysine 120g, BHT, 5g. HSM = Heat-treated Sheep Manure, SBM = Soyabean Meal, PKC = Palm Kernel Cake and E = Enzyme.

Table-3. Effect of Roxazyme supplementation of heat-treated sheep manure on the performance of the birds.

	T ₁	T ₂	T ₃	T ₄	T ₅	T ₆	T ₇	
% Heat- treated sheep manure and roxazyme G inclusion								
Performance criteria	0%	5%	5%+E	10%	10%+E	15%	15%+E	SEM
Av. Initial body weight (g)	63.16	62.48	63.11	62.94	62.92	62.88	63.09	
A.V Final body weight (g)	680.44 ^{bc}	665.11 ^{bc}	761.01 ^a	650.68 ^c	703.42 ^b	591.87 ^d	623.12 ^{bc}	4.19
Av. Body weight gain (g)	617.28 ^{bc}	602.63 ^{bc}	697.90 ^a	587.74 ^c	640.50 ^b	528.98 ^d	623.12 ^{bc}	4.19
Av. Daily weight gain (g)	17.64 ^{bc}	17.22 ^{bc}	19.94 ^a	16.79 ^c	18.30 ^b	15.11 ^d	17.80 ^{bc}	0.17
Av. total feed intake (g)	1732.03 ^d	1789.32 ^{cd}	1739.62 ^{cd}	1869.70 ^b	1789.69 ^{cd}	1978.90 ^a	1829.22 ^{bcd}	10.74
Av. Daily feed intake (g)	49.48 ^d	51.12 ^{bcd}	49.70 ^{cd}	53.4 ^{2b}	51.13 ^{cd}	56.54 ^a	52.26 ^{bc}	0.93
Feed conversion ratio	2.81 ^b	2.97 ^b	2.50 ^a	3.18 ^c	2.80 ^b	3.74 ^d	2.94 ^b	0.35
Av. Daily protein intake (g)	10.75 ^{ab}	10.85 ^{ab}	10.55 ^a	11.29 ^{bc}	10.80 ^{ab}	11.90 ^{bc}	11.00 ^{ab}	0.41
Protein efficiency ratio	1.64 ^b	1.59 ^{bc}	1.84 ^a	1.49 ^c	1.69 ^b	1.27 ^d	1.62 ^b	0.25

^{a,b,c,d,e} Means with different superscripts on same row differ significantly (P < 0.05)



CONCLUSIONS

Results of this study indicated that body weight gain, feed intake, feed conversion ratio and protein efficiency ratio of starter broilers were negatively affected by 15% unsupplemented HSM dietary inclusion level. Enzyme supplementation of HSM broiler starter diets enhanced the performance, reduced the cost of feed intake and cost of feed per weight gain of the of the young broiler chicks. It may be concluded that that enzyme supplementation of higher level of HSM might be effective for better performance of the birds.

REFERENCES

- A.O.A.C. 1995. Official Methods of Analysis (6th Ed) Association of Analytical Chemist. Washington D.C.
- F.O. Abeke, S. O. Ogundipe, I.A. Adeyina, B.Y. Abubakar, O.O. Oni and B.I. Nwagu. 2003. Response of laying hens to dietary levels of heat-treated sheep manure (HSM). *Tropical Journal of Animal Science*. 6(2): 111-116.
- F.O. Abeke, S. O. Ogundipe, A. A. Sekoni, J. O. Bale, I. A. Adeyinka and O. O. Oni. 2008. Effects of dietary levels of heat - treated sheep manure (HSM) on the performance and egg quality parameters of laying hens. *Nigerian Journal of Animal Production*. 35(1): 56-62.
- T. Acamovic. 2001. Commercial application of enzyme technology for poultry production. *World's Poultry Science Journal*. 57: 225-242.
- A. O. Aduku. 1993. Tropical Feedstuff Analysis Table. Department of Animals Science, Faculty of Agriculture Ahmadu Bello University, Zaria.
- K. Ajaja, J.O. Agbede and V.A. Aletor. 2003. Influence of roxazyme supplementation on the utilization of wheat offal or rice by broilers. *Proceedings of the 8th Annual Conference Animals Science Association of Nigeria*, Minna. pp. 32-34.
- R.A. Alabi and M.R. Aruna. 2005. Technical efficiency of family poultry production in Niger Delta, Nigeria. *Journal of Central European Agriculture*. 6(4): 531-538.
- O. A. Alalade and E. A. Iyayi. 2006. Chemical composition and the feeding value of Azolla (*Azolla pinnata*) meal for egg - type chicks. *International Journal of Poultry Science*. (2): 137-141.
- W. An. 1994. Effects of different sources of fibre on Performance of tibia mineral content in broilers. *CAB International Poultry Abstract*. 20: 377
- A. O. Ani and O.D. Omeje. 2007. Effect of supplementation with enzyme on growth performance of broiler chicks fed diets containing raw bambara nut (*Voandoeia Subterranean*) Waste. *Proceedings of the 32nd Annual Conference of the Nigerian Society for Animal Production*, Calabar, Nigeria. pp. 278-281.
- M. R. Bedford. 1997. Reduced viscosity of intestinal digesta and enhanced nutrient digestibility in chicken given exogenous enzymes. In: *Enzymes in poultry and swine nutrition*. IDRC Publication. p. 145.
- Federation of Animal Science Societies (FASS). 1999. Guide for the care and management of agricultural animals in agricultural research and teaching. 1st revised Ed. Savoy, IL. p. 58
- M. Gunal. and S. Yasar. 2004. Performance and some digesta parameters of broiler chickens given low or high viscosity wheat based diets with or without enzyme supplementation. *Turkish Journal Veterinary and Animal Science*. 28: 223-327.
- M.E. Jackson. K. Geronian, A. Knox, J. McNab and E. McCartney. 2004. A dose-response study with the feed enzyme β -mannanase in broilers provided with corn-soybean meal based diets in the absence of antibiotic. *Poultry Science*. 83: 1992-1996.
- W. Kongbuntad, C. Khanongnuch and S. Lumyong. 2006. Efficacy of zylanase supplementation produced from *Thermoascus aurantiacus* SL 16w in diet on Thai native chicken performance. *International Journal of Poultry Science*. 5(5): 463-469.
- National Research Council. 1994. *Nutrient Requirement of Poultry*, 9th revised edition. National Academy of Science, Washington, D.C
- I. U. Obi. 2002. Statistical methods of detecting differences between treatment means and research methodology issue in laboratory and field experiments. A.P Company Ltd.
- P. M. P. Oldalle and F. Hoffman. 1996. Enzymes -a tool for unlocking nutrients in animal feeds Roche Nigeria Seminar, Lagos, Nigeria. pp. 56-62.
- P. N. Onu. 2007. The influence of heat- treated sheep manure on the performance, carcass characteristics and economics of production of starter broilers. *Journal of Animal and Veterinary Advances*. 6(11): 1323-1327.
- P. N. Onu and M. O. Otuma. 2008. Utilization of heat - treated sheep chopping in the diets of broiler finisher chicks. *International Journal of Poultry Science*. 7(2): 169-173.
- P. N. Onu. 2006. Performance response of starter broilers fed Ronozyme in soyabean hulls-based diets. *Journal of Science, Engineering and Technology*. 13(1): 6664-6674.



- P. N. Onu, P. E. Nwakpu and L.O. Chukwu. 2006. Performance of broiler chicks (*Gallus domesticus*) fed maize offal-based diets supplemented with roxazyme G. enzyme. International Journal of Poultry Science. 5(7): 610.
- G. Richter, A. Lemser and A. Muller. 1995. Is enzyme feeding beneficial in broiler feeding? Lohnt Sich der Einsat, Von enzyme in der Bralermast? Muhle + Mischfutter technik. 131(39): 546-547. In: Poultry Abstract. 21: 5.
- A. A. Saki, M.T. Mazugi and A. Kamyab. 2005. Effect of mannanase on broiler performance, ileal and *invitro* protein digestibility, uric acid and litter moisture in broiler feeding. International Journal of Poultry Science. 4: 21-26.
- K. Samarasinghe, R. Messikommer and C. Wank. 2000. Supplementation enzymes and their effect on nutrient utilization and growth performance of growing chickens as affected by pelleting temperature. Archives of Animals Nutrition. 53: 45-58.
- A. Teimour, M. Rezae, J. Pourreza, H Sayyazhedch and P. W. Waldroup. 2005. Effect of diet dilution in the starter period on the performance and carcass characteristics of broiler chicks. International Journal of Poultry Science. 4(12): 1006-1011
- V. Tufarelli, M. Dario and V. Laudadio. 2007. Effect of xylanase supplementation and particle-size on performance of guinea fowl broilers fed wheat based diets. International Journal of Poultry Science. 4: 302-307.
- P. J. Van Soest, J.B. Robertson and B.N. Lewis. 1991. Food and Fibre. Marabon Symposium Navings Parsking supple. 14: 12.
- S. M. Yashim, F. Onya, S. B. Abdu, H. Y. Adamu and M. A. Dakare. 2007. Castor (*Ricinus communis* Linn.) fruits as a source of livestock feed. Proceedings of the 32nd Annual Conference of the Nigerian Society for Animal Production, Calabar. pp. 336-337.
- B. I. Zobell, M Choct and R. D. Freed. 2000. Intuition of nutrient digestion by wheat Pentosans. British Journal of Nutrition. 67: 123-132.
- X. T. Zou, X.J. Qiao and Z.R. Xu. 2006. Effect of β -mannanase (Hemicell) on growth performance and immunity of broilers. Poultry Science. 85: 2176-2182.
- H. A. H. Zakaria, A. R. J. Mohammad and S. J. Amer. 2008. Effect of exogenous enzymes on the growing performance of broiler chickens fed regular corn/soybean-based diets and the economics of enzyme supplementation. Pakistan Journal of Nutrition. 7(4): 534-539.