

www.arpnjournals.com

# CROPPING SYSTEM ANALYSIS USING REMOTE SENSING AND GIS: A BLOCK LEVEL STUDY OF KURUKSHETRA DISTRICT

M. P. Sharma<sup>1</sup>, Manoj Yadav<sup>1</sup>, R. Prawasi<sup>1</sup>, Pavan Kumar<sup>2</sup> and R. S. Hooda<sup>1</sup> <sup>1</sup>Haryana Space Applications Centre (Department of Science and Technology), Hisar, India <sup>2</sup>Department of Remote Sensing, Banasthali University, Rajsthan, India E-Mail: <u>pawan2607@gmx.com</u>

# ABSTRACT

Agricultural sustainability has the highest priority in all countries, whether developed or developing. Cropping System Analysis is essential for studying the sustainability of agriculture. Crop rotation is stated as growing one crop after another on the same piece of land in different timings (seasons) without impairing the soil fertility. A cropping system can be defined as the cropping patterns and their management to derive maximum benefits from a given resource base under specific environmental conditions. Multiplicity of cropping system has been one of the main feature of Indian agriculture and is attributed to rained agriculture and prevailing socio-economic situations of farming community. It has been estimated that more than 250 double cropping systems are followed throughout the country. Thirty important cropping systems have been identified based on rationale spread of crops in each district in the country. Although, it is well known that one of the main advantage of remote sensing satellites is the synoptic and repeated collection of data which facilitate to map multi-year cropping patterns and crop rotations. In the present work, crop rotation and long term changes monitoring in cropping pattern along other spatial and non-spatial collateral data have been done with the help of satellite data at block level of Kurukshetra district of Haryana. Multi-date IRS LISS-III data of different seasons for the year 2007-08 have been used for the study. Cropping pattern maps of Rabi, Kharif and Summer season have been understood to know the spatial distribution and associations between crops or crops and uncultivated land in the same fields (although not in a particular order of sequence). The findings of the study may be used by Department of Agriculture, Haryana for planning of agricultural strategies in the district and for planning agricultural research and extension activities for crop diversification.

Keywords: cropping system, remote sensing, IRS-P6 satellite, LISS-III, Rabi, Kharif.

### **1. INTRODUCTION**

Agricultural sustainability has the highest priority in all countries, whether developed or developing. Cropping System Analysis is essential for studying the sustainability of Agriculture. Indian society is agriculture based and its economy is dependent on agriculture. Since the scope of increasing area under agriculture is rather limited thus, to increase agriculture productivity. Majority of Indian farmers get the major share of income from crop production. Therefore, it is very much important to select the right crop, in the right season so that maximum profit may be achieved. In selecting a crop for a season, both post and pre season crops should also to be examined (Manjunath K. R. 2006). In agricultural applications, remote sensing imagery has been used to identify different crop types, estimate crop area and, predict yield at small scales (Kanemasu, 1974). The structure of most crops is identical causing spectral mixes within crops and other types of vegetation. These high spectral overlaps make attempts to understand the relationships between crops and the ecosystems within which they occur in order to classify remote sensing imagery difficult. The spectral signature for vegetation is highly variable in nature since it changes completely during the seasonal cycle of many plants (Schowengerdt, 1997). Therefore, a number of contexts like spatial, spectral and temporal have been used over the years in order to carry out crop identification in remotely sensed data (Byeungwoo and Landgrebe, 1992; Wharton, 1982).

Agriculture is the backbone of Indian economy, contributing about 90% towards the Gross National product and providing livelihood to about 70% of the population. In the country, wheat is the second important food crop being next to rice and contributes to the total food grain production of the country about 25%. Importance of crop production information was realized in India as early as 1884, when the government initiated a program for wheat. Agriculture is the main occupation of the people of Haryana. Haryana is often called the "Food Mine" of the country. About 80% of the population of the state is agriculture dependent. Haryana is self sufficient in producing food grains of the country. Wheat, rice, maize and bajra are the major grains produced in the state. The crop production of Haryana can be broadly divided into Rabi and Kharif. The major Kharif crops of Haryana are rice, jowar, bajra, maize, mustard, jute, sugarcane, sesame and groundnut. The major Rabi crops are wheat, tobacco, gram, linseed, rapeseed and mustard. The total area irrigated by canal water is 21.40 lakh hectares. The state of Haryana has a geographical area of 44.20 lakh hectare. About 86% of the geographical area is cultivable in which 96% has already been brought under plough. Therefore, there is hardly any scope for bringing additional area under cultivation, except for reclamation of degraded lands affected by water logging, salinity and alkalinity. About 84% of the cultivated area in the state is irrigated. Irrigation from canals forms the lifeline of agriculture in Haryana. The various canals which are operating in the state include Western Yamuna Canal, Gurgaon Canal, Jui



## www.arpnjournals.com

Canal, Jawaharlal Lal Nehru Canal and Bhakra Canal. These canals are the main source of water for cultivation in various districts of the state. Haryana together with Punjab is called the 'Grain Bowl' of India.

### Objective

This study established the methodology for spatial analysis of cropping systems. It is now envisaged to create remote sensing based Cropping System Analysis for Kurukshetra district and its development Blocks at 24 m. cell size.

### 2. STUDY AREA

The Imperial Gazetteer of India (Vol. XIII) records (Imperial Gazetteer of India, Vol. III) that the word Haryana is probably derived from Harialban or the land of green forests and is reminiscent of the ancient time when this area was a rich and fertile tract (F. Wilson, 1904). Also subscribes to this view in 'Punjab Notes and Queries'. Kurukshetra district falls in the north-east part of the Haryana State and is bounded by North latitudes 29°53'00" and 30°15'02" and East longitudes 76°26'27" and 77°07'57". It falls in parts of Survey of India Toposheets No. 53B and 53C covering an area of 1530 sq. km. The district covers 3.46% area of the State. The district is bordered by Karnal district in the south, Kaithal district in the south and south-west, Ambala in the north and Patiala in the north-west. The plain is remarkably flat and narrow low-lying flood plains, known as either Betre Khadar of Naili. Saraswati, Markanda and Ghaggar rivers are major rivers of the district. The eastern parts of the district falls in the Upper Jamuna Basin and western parts falls in Ghaggar basin. The river Markanda provides the major drainage in the area. Irrigation in the district is done by surface water as well as ground water.

### 2.1 Development blocks of Kurukshetra district

There are five development blocks in Kurukshetra district.



Figure-1. Map of study area.

#### 2.1.1 Pehowa block

Pehowa Block is one of the development blocks of Kurukshetra district which has location  $29^{0}52'30''-30^{0}6'3''N$  and  $76^{\circ}27'00''-76^{\circ}50'00''E$ . The total geographical area of the Pehowa Block is 500.17 sq. km.

### 2.1.2 Shahabad block

The Block lies along the north border of the Kurukshetra district between  $30^{\circ}03'30''$  to  $30^{\circ}15'30''$  North latitudes and  $76^{0}38'30''$  to  $76^{0}58'30''$  East longitudes in Kurukshetra district. Total geographical area of this Block

is 382.72 sq. km. The area being on a river bank is fertile and is used for paddy as well as wheat and vegetables.

#### 2.1.3 Thanesar block

The location of Thanesar Block in Kurukshetra district is between  $29^{\circ}52'00''$  to  $30^{\circ}07'30''$ North and  $76^{\circ}40'30''$  to  $77^{\circ}00'00''$  East. Total geographical area is 517.11 sq. km.

#### 2.1.4 Ladwa block

#### www.arpnjournals.com

© 2006-2011 Asian Research Publishing Network (ARPN). All rights reserved.

This Block is situated between  $29^{\circ}56'30'' - 30^{\circ}04'40''$  North latitudes and  $76^{\circ}55'00'' - 77^{\circ}07'00''$  East longitudes. Total geographical area of this block is 170.44 sq. km.

### 2.1.5 Babain block

The location of Babain Block in Kurukshetra district is between  $30^010'00"$  to  $30^\circ00'00"$  North and  $76^\circ55'30"$  to  $77^\circ05'30"$  East. Total geographical area is 172.58 sq. km. Surrounding Blocks of Babain are Ladwa Block in south, Shahabad Block in north-west and Thanesar Block in south-west. It is covered by Yamunanagar district in north-east and Ambala district in north.

### **3. DATABASE REQUIREMENTS**

#### 3.1 Remote sensing data

Remote sensing data is the basic data source for mapping the cropping system of the state region. Indian Remote Sensing Satellite Resourcesat (IRS-P6) LISS III data is the ideal one with optimum spatial and temporal resolution. The sensor provides 23.5 m spatial resolution data in Green, Red, NIR and SWIR bands with 24 days revisit capability. Its repeat cycle can be used for deriving Kharif, Rabi and summer cropping pattern and change analysis between these seasons. Remote sensing data from sensor LISS III on-board Indian Remote Sensing Satellite Resourcesat (IRS-P6) of 2007-08 were used to analyze the changes in cropping pattern and crop rotation for three seasons' Kharif, Rabi and summer. The Multi-date satellite data are used for different seasons which are given in Table-1.

Table-1.	Satellite	data	used 1	in digital	analysis.

.. . .

S. No.	Season	Satellite	Sensor Date of acquisition		Path/Row
				October 24, 07	
1.	Kharif	IRS-P6	LISS-III	July 15, 07	95/50
				Sept. 30, 07	
2	Rahi	IDS D6	1 155 111	Jan. 4, 08	95/50
۷.	Kabi IN	113-10	L155-111	March 16, 08	95/50
3.	Summer	IRS-P6	LISS-III	May 27, 08	95/50



Figure-2. Flow chart of methodology used.

#### 3.2 Cropping system analysis



#### www.arpnjournals.com

The IRS-LIII data was the main source of satellite imageries for deriving the spatial maps of agricultural land use, cropping pattern, crop rotation and crop calendar. Total-6 dates starting from September 30, 2007, October 24, 2007, July 15, 2007, January 4, 2008, March 16, 2008 and May 27, 2008 were used. PCI Geomatica v9.3 image processing software was used for geo-referencing, multi-date data stacking and overlaying state and district boundary. The best channel selection was carried out with all the dates methodology used that resulted in 6 optimum dates. Optimum data sets were used to classify forest/permanent vegetation, wastelands and permanent water bodies. A multi-phased unsupervised ISODATA classification was used for seasonal cropping pattern mapping. The pixels that were found as correctly classified were masked and a second classification was performed for remaining area until no further increase in class separability was achieved. Data acquired from September 30, 2007, October 24, 2007, July 15 and January 4, 2008, March 16, 2008 and May 27, 2008 or Kharif, Rabi and summer cropping patterns, respectively.

# 4. RESULTS

Analysis by remote sensing techniques reflected that Rice and Sugarcane are two major crops in Kharif season, which could be identified using Multi-date RS data. Rice is mostly concentrated throughout the district except in eastern part of the district. Other crops in the Kharif season include Bajra, Maize, Jowar etc. Wheat and Sugarcane are the two major crops during Rabi season followed by other crops.

# 4.1 Cropping system analysis

Crop rotation shows sequential planting of crops in time. It indicates that Kurukshetra District has the major crop rotations of Rice-Wheat-other Crops, Rice- Wheat-Fallow, Fallow-Wheat-Fallow and Sugarcane based of three seasons Kharif, Rabi and summer. Rice-Wheat-other Crops, Rice- Wheat-Fallow, Fallow-Wheat-Fallow and Sugarcane based rotation is distributed in whole district except eastern part of the district shown in Map No. 4. This may be due to the effect of Rice and Wheat, major crops of Kharif and Rabi season, are gradually spreading from north to south of Kurukshetra District. In summer season, fewer crops are grown and most of the agriculture area is Fallow. So the Kurukshetra district has major agricultural area of double crop and less area of more than two crops. The presence of other crops is slowly getting reduced, especially in Rabi Season.

# 4.1.1 Pehowa block

It is indicated that Pehowa Block has major crop rotations of Rice-Wheat-Fallow, Rice-Other Crop-Fallow, Fallow-Wheat Fallow and Sugarcane based of three seasons as can be seen in Table-2 and Figure-3. Rice-Wheat-Fallow rotation dominates in the western and northern part of the district while Sugarcane Based rotation dominates in the eastern part of the block. This may be due to the effect of Rice and Wheat, major crops of Kharif and Rabi season respectively, and due to Fallow in summer. Rice-Wheat-Fallow rotation means that the part of land where Rice grows in Kharif season, Wheat in Rabi season and Fallow in summer season. Sugarcane based is also the major rotation in this block.

Table-2. Crop rotations statistics of Pehowa block.

S. No.	Rotations	<b>Area</b> (000'h)	
1.	Rice-wheat-sunflower	0.30	
2.	Rice-wheat-saathi	0.02	
3.	Rice-wheat-other crops	6.02	
4.	Rice-wheat-fallow	25.82	
5.	Fallow-wheat-other crops	0.87	
6.	Other crops-wheat-other crops	1.51	
7.	Fallow-wheat-sunflower	0.06	
8.	Fallow-wheat-fallow	2.32	
9.	Fallow-fallow-other crops	0.21	
10.	Rice-fallow-fallow	0.27	
11.	Fallow-fallow-sunflower	0.02	
12.	Fallow-fallow-fallow	0.36	
13.	Sugarcane based	1.13	
14.	Other rotation	6.36	



Figure-3. Crop rotations map of pehowa block.

# 4.1.2 Thanesar block



www.arpnjournals.com

Thanesar Block has the major crop rotations of Rice-Wheat-Fallow of three seasons as it can be seen in Table-3 and Figure-4. Rice-wheat-Fallow rotation also dominates in the southern part of the district while Sugarcane Based rotation dominates in the some middle and eastern part of this block.

VOL. 6, NO. 10, OCTOBER 2011

C N-	Detetions	A (000)1			
Table-3. Crop rotations statistics of Thanesar block.					

S. No.	Rotations	<b>Area</b> (000'h)
1.	Rice-wheat-sunflower	0.52
2.	Rice-wheat-saathi	0.09
3.	Rice-wheat-other crops	6.83
4.	Rice-wheat-fallow	18.85
5.	Fallow-wheat-other crops	1.64
6.	Other crops-wheat-other crops	1.41
7.	Fallow-wheat-sunflower	0.01
8.	Fallow-wheat-fallow	3.30
9.	Fallow-fallow-other crops	0.50
10.	Rice-fallow-fallow	0.55
11.	Fallow-fallow-sunflower	0.06
12.	Fallow-fallow-fallow	0.69
13.	Sugarcane based	3.36
14.	Other rotation	8.08



Figure-4. Crop rotations map of Thanesar block.

# 4.1.3 Shahabad block

Shahabad Block has the major crop rotations of Rice-Wheat-Other Crops of three seasons as can be seen in Table-4 and Figure-5. Rice-wheat-Other Crops rotation also dominates in the western and northern part of the district while Sugarcane Based rotation dominates in the some eastern and southern part of this block Out of total agriculture area 7.95 (000'ha) area of rotations are comes under other rotations.

Table-4. Crop rotations statistics of Shahabad block.

S. No.	Rotations	Area (000'h)
1.	Rice-wheat-sunflower	1.51
2.	Rice-wheat-saathi	0.01
3.	Rice-wheat-other crops	8.56
4.	Rice-wheat-fallow	6.78
5.	Fallow-wheat-other crops	1.39
6.	Other crops-wheat-other crops	0.52
7.	Fallow-wheat-sunflower	0.27
8.	Fallow-wheat-fallow	0.99
9.	Fallow-fallow-other crops	0.79
10.	Rice-fallow-fallow	0.53
11.	Fallow-fallow-sunflower	0.20
12.	Fallow-fallow-fallow	0.28
13.	Sugarcane based	5.59
14.	Other rotation	7.95



Figure-5. Crop rotations map of Shahabad block. 4.1.4 Babain block

# www.arpnjournals.com

Babain Block has the major crop rotations of Rice Wheat-Other Crops of three seasons as can be seen in Table-5 and Figure-6. Rice-Wheat-Other Crops rotation also dominates in the western, southern and northern part of the district while Sugarcane based rotation dominates in some northern and southern part of this block. Out of total agriculture area 3.44 (000'ha) area of rotations are comes under other rotations.

S. No.	Rotations	<b>Area</b> (000'h)
1.	Rice-wheat-sunflower	0.72
2.	Rice-wheat-saathi	0.00
3.	Rice-wheat-other crops	3.97
4.	Rice-wheat-fallow	1.58
5.	Fallow-wheat-other crops	0.55
6.	Other crops-wheat-other crops	0.33
7.	Fallow-wheat-sunflower	0.08
8.	Fallow-wheat-fallow	0.25
9.	Fallow-fallow-other crops	0.19
10.	Rice-fallow-fallow	0.12
11.	Fallow-fallow-sunflower	0.03
12.	Fallow-fallow-fallow	0.06
13.	Sugarcane based	3.88
14.	Other rotation	3.44





Figure-6. Crop rotations map of Badain block.

The Block has two major crop rotations: Rice-Wheat-Other crops and sugarcane based. The crop rotations Rice-Other crops-Fallow and Sugarcane based occupies 3.35 and 3.85 (000'ha) area respectively. Rice-Wheat-Other crops rotation also dominates in the northern and southern part of the district while Sugarcane based rotation dominates in some eastern and southern part of this block as can be seen in Table-6 and Figure-7, shows the area of different crop rotations in the block. Out of total agricultural area 3.22 ('000 ha) area comes under other rotations.

S. No.	Rotations	<b>Area</b> (000'h)
1.	Rice-wheat-sunflower	0.64
2.	Rice-wheat-saathi	0.03
3.	Rice-wheat-other crops	3.35
4.	Rice-wheat-fallow	1.64
5.	Fallow-wheat-other crops	0.52
6.	Other crops-wheat-other crops	0.47
7.	Fallow-wheat-sunflower	0.08
8.	Fallow-wheat-fallow	0.28
9.	Fallow-fallow-other crops	0.17
10.	Rice-fallow-fallow	0.12
11.	Fallow-fallow-sunflower	0.03
12.	Fallow-fallow-fallow	0.11
13.	Sugarcane based	3.85
14.	Other rotation	3.22

Table-6. Crop r	rotations	statistics	of I	Ladwa	block
-----------------	-----------	------------	------	-------	-------

ARPN Journal of Agricultural and Biological Science

 $\ensuremath{\textcircled{\sc 0}}$  2006-2011 Asian Research Publishing Network (ARPN). All rights reserved.



#### www.arpnjournals.com



VOL. 6, NO. 10, OCTOBER 2011

Figure-7. Crop rotations map of ladwa block.

# 5. CONCLUSIONS

The present study introduces a method for analysis of cropping systems for main crops, other crops and fallow land of the year 2007-08 of the Kurukshetra district and its development blocks. Using unsupervised classification of multi-date satellite images of LISS III, cropping pattern and crop rotation maps are generated. This method identifies main crops of Kharif, Rabi and Summer Seasons shown in the study area. Although, it shows crop rotations as such, it allows spatial relationships between main crops, and other crops at a specific location to be mapped during the study period. An advantage of the proposed method is that it may show some spatial relationships between crops, which could reveal certain specific location of some rotations. For example, in the study area of Kurukshetra district the spatial relationship between Rice, Wheat and Sunflower which are seasonal crop in Kharif, Rabi and summer shows the specific location of typical distribution areas. An additional possibility of multi-year cropping pattern map is its use in future spatial crop distribution prediction, since it contains expert knowledge about spatial relationships between crops in the study area and implicit probabilities of changes.

# REFERENCES

Manjunath K.R. 2006. Remote Sensing and GIS Applications for Crop Systems Analysis. Invited lecture delivered during NNRMS Training Programme on "Geoinformatics for Sustainable Development" at, Haryana Remote Sensing an applications Centre, Hisar, india.

Panigrahy Sushma. Ray S.S. Sood Anil. Patel L.B. Sharma P.K. and Parihar J.S. 2004. Analysis of Cropping Pattern changes in Bhatinda District Punjab. Journal of the Indian Society of Remote Sensing. 32(2): 209-216.

Panigrahy R. K. Panigrahy S. Ray S. S. and Parihar J. S. Analysis of the Cropping pattern and Crop rotation of a subsistence Agricultural region using high temporal Remote Sensing data.

Saha S.K. and Choudhury Swagata. Cropping Pattern Change Analysis and Optimal Landuse Planning By Integrated Use Of Satellite Remote Sensing and GIS - A Case Study Of Barwala C.D. Block, Panchkula District, Haryana, India.

Imperial Gazetteer of India. 13: 54(3).

Wilson F. 1904. Punjab Notes and Quarries. 547: 67.

Byeungwoo J. and Landgrebe D. A. 1992. Classification with spatio - temporal interpixel class dependency contexts. IEEE Transactions on geosciences and remote sensing. 30(4): 663-672.

Kanemasu E. T. 1974. Seasonal canopy reflectance patterns of wheat sorghum and soybean. Remote Sensing of Environment. 3(1-4).

Schulze R. E. 1997. South African Atlas of Agrohydrology and Climatology (Report TT82/96). Pretoria: Water Research Commission.

Wharton S. W. 1982. A contextual classification method for recognizing land use patterns in high resolution remotely sensed data. Pattern Recognition. 1(4): 317-324.