



## STUDY EFFECT OF IBA HORMONE LEVELS ON ROOTING IN MICRO CUTTINGS OF TEA (*Camellia sinensis* L.)

Sirous Bidarigh<sup>1</sup> and Ebrahim Azarpour<sup>2</sup>

<sup>1</sup>Department of Agriculture, Lahijan Branch, Islamic Azad University, Lahijan, Iran

<sup>2</sup>Young Researchers Club, Lahijan Branch, Islamic Azad University, Lahijan, Iran

E-Mail: [Bidarig@yahoo.com](mailto:Bidarig@yahoo.com)

### ABSTRACT

There has been a steady demand for production and supply of cuttings of different varieties for tea gardens in Iran. In order to study the effect of indole-3-butyric acid (IBA) hormone levels (0, 0.1; 0.5, 1 and 3 mg/L) on rooting in micro cuttings of tea (*Camellia sinensis* L.), an experiment as RCBD with four replications was conducted during 2010 at Research Laboratory of Faculty of Agriculture, Lahijan University in Iran. The culture bed was contained MS, sucrose (3%) and agar (75%). The results show that IBA hormone levels on root length and root number in micro cuttings of tea had a significant difference in 1% probability level. The highest root length (17.22 mm) and root number (4.93) in micro cuttings of tea were obtained with application 1mg/L IBA. The lowest root length (3.17 mm) and root number (0.68) in micro cuttings of tea were obtained with application 0mg/L IBA.

**Keywords:** tea, micro cutting, IBA hormone, root length, root number.

### INTRODUCTION

Single leaf-node cuttings conventionally propagate cultivated tea. This method of production becomes limiting when large numbers are required from new clones of which only very few stock plants are available. Moreover, the process in establishing tea plants from cutting is lengthy and labour-intensive, rendering it ineffective for rapid dissemination of new clones for large-scale commercial plantings (Tahardi *et al.*, 2003). Conventional tea breeding is well established, though time-consuming and labor intensive due to its perennial nature and long gestation period (4-5 years). Additionally, tea breeding has been slowed by lack of reliable selection criteria. Vegetative propagation is standard, yet limited by slow multiplication rate, poor survivability of some clones, and need for copious initial planting material (Tahardi *et al.*, 2003). Seed-borne plants are heterogeneous due to their highly allogamous nature; consequently, it is difficult to maintain their superior character. Research on tea micro-propagation has recently focused on exploring the potential of somatic embryogenesis as a more efficient means of plant manipulation and regeneration (Tahardi *et al.*, 2003). Somatic embryos have been used as a model system to understand the mechanisms regulating plant embryogenesis, being an alternative for the propagation of plants with high rates of multiplication, with relevance in tree improvement programmes (Sghaier *et al.*, 2009). To induce calli from tea explants and to cultivate the calli to produce plants, various kinds of media have been designed. Generally, it would be better to use only one or two kinds of basal media in combination of different kinds and concentrations of phytohormones (Misawa, 1994). The most suitable medium composition should be optimized afterwards in order to obtain higher level of products as well as higher growth rate. Plant hormones play essential roles in plant metabolism and can influence cell cycle proteins (Sánchez *et al.*, 2005). Tea (*Camellia*

*sinensis* L.) is the oldest non-alcoholic caffeine-containing beverage crop in the world and health benefits attributed to tea consumption are well proven (Mondal, 2004). Tea is one of the most important plantation crops in the world. At present, the most parts of tea fields were planted with various clonal propagated cultivars. Fields performance of micro-propagated tea plants and the impact of cultural operations on growing tea plants have been reported previously (Marimuthu and Raj Kumar, 2001). For these reasons, tea plants were introduced into tissue culture for plant regeneration and genetic manipulation. Unlike other crops, reports are not available on the basic physiology of micro-propagated tea plants Marimuthu and Raj Kumar, 2001).

In vitro culture is one of the key tools of plant biotechnology that exploits the totipotency nature of plant cells, a concept proposed by Haberlandt (1902) and unequivocally demonstrated, for the first time, by Steward *et al.* (1985). Tissue culture is alternatively called cell, tissue and organ culture through in vitro condition (Debergh and Read, 1991). It can be employed for large-scale propagation of disease free clones and gene pool conservation. Ornamental industry has applied immensely in vitro propagation approach for large-scale plant multiplication of elite superior varieties. As a result, hundreds of plant tissue culture laboratories have come up worldwide, especially in the developing countries due to cheap labour costs. However, micropropagation technology is more costly than conventional propagation methods, and unit cost per plant becomes unaffordable compelling to adopt strategies to cut down the production cost for lowering the cost per plant (IAEA-TECDOC, 2004).

Rooting the shoots produced in vitro or micro cuttings, has been achieved through in vitro and ex vitro or non-sterile, conditions (Briggs and McCulloch, 1984). In some cases, micro cuttings root better in vitro



environments. In vitro rooting was superior to ex vitro rooting for *Prunus* x 'Hally Jolivette' (Lineberger, 1983). Also, in some cases in vitro, it is beneficial to make changes to the medium. Li and Eaton (1984) reported that rooting 'Marechal Foch' grapevine in half-strength MS salts was superior to rooting in full strength MS salts. But in other cases, superior rooting can result under ex vitro conditions. Rooting of *Halesia carolina* L. non-sterile conditions was reported to be far superior to rooting in sterile conditions (Brand and Lineberger, 1986).

The objective of the present research was to enhance the development of a rooting in micro cuttings of tea (*Camellia sinensis* L.) with application IBA hormone levels.

## MATERIALS AND METHODS

In order to study the effects of indole-3-butyric acid (IBA) hormone levels (0, 0.1; 0.5, 1 and 3 mg/L) of in vitro culture of tea (*Camellia sinensis* L.), an experiment as RCBD with four replications was conducted during 2010 at Research Laboratory of Faculty of Agriculture, Lahijan University in Iran. Tea apex meristems were obtained from 2-years-old rooted cuttings of tea plants grown in a green house, with a 16 h photoperiod and photosynthetic photon flux of  $101.5 \mu\text{mol m}^{-2} \text{S}^{-1}$  (400-700 nm) at the plant level. Leaf samples were washed with tap water and surface sterilized in a drop of liquid detergent for 1 min, followed by three rinses in sterile distilled water. Then, they were re-sterilized with 70% ethanol for 30s and with 20% sodium hypochlorite for 20 min, followed by three rinses in sterile distilled water, all under laminar flow. Discs of ca. 0.5 cm<sup>2</sup> diameter were cut from the leaves and were cultured on MS (Murashige and Skoog, 1962) basal medium (Figure-1). PH was adjusted to 5.8 before adding 3% (w/v) sucrose and 75% (w/v) agar. Autoclaving was done for 20 min at 120°C and 150 KPa. The cultures were maintained with a 16 h photoperiod at  $27 \pm 3^\circ\text{C}$ . For the induction of callus to rhizogenesis, the calli derived from micro cuttings were cultured on application indole-3-butyric acid (IBA) hormone levels (0, 0.1; 0.5, 1 and 3 mg/L) only.

Data analyses were analyzed by using SAS software. The Duncan's multiple range tests was used to compare the means at 5% of significant.

## RESULTS AND DISCUSSIONS

Results of variation analysis show that (Table-1), IBA hormone levels on root length in micro cuttings of tea had a significant difference in 1% probability level. The highest root length (17.22 mm) in micro cuttings of tea was obtained with application 1mg/L IBA and the lowest root length (3.17 mm) was obtained with application 0mg/L IBA (Figure-2). Results of variation analysis show that (Table-1), IBA hormone levels on root number in micro cuttings of tea had a significant difference in 1% probability level. The highest root number (4.93) in micro cuttings of tea was obtained with application 1mg/L IBA and the lowest root number (0.68) was obtained with application 0mg/L IBA (Figure-3). Result of analysis of

liner (Figure-4) show that between root length and root number a positive and very significant correlation ( $R^2 = 0.99$ ).


Micro propagation generally involves four distinct stages: initiation of cultures, shoot multiplication, rooting of in vitro grown shoots, and acclimatization. The first stage: culture initiation depends on explant type or the physiological stage of the donor plant at the time of excision. Explants from actively growing shoots generally used for mass scale multiplication. The second stage: shoot multiplication is crucial and achieved by using Plant Growth Regulators i.e., auxin and cytokinin. The third stage: the elongated shoots, derived from the multiplication stage, are subsequently rooted either ex vitro or in vitro. In some cases, the highest root induction occurs from excised shoots in the liquid medium when compared with semi-solid medium. The fourth stage: acclimatization of in vitro grown plants is an important step in micro propagation (Smart, 2008; Rout *et al.*, 2006). This greening process was also observed in our experiments. The effect of abscisic acid (ABA) and excess benzyladenine (BA) on the formation of shoot from tea (*Camellia sinensis* L.) leaf was investigated (Ghanatia and Rahmati Ishkaa, 2009); As a result, we observed root formation in the transferred calli to B5 medium after one month. Subsequently, the calli were transferred to the aforesaid B5 medium supplemented with emergence of root.

ABA (2 mg/L) and excess BA (400 mg/L) to form shoot. The calli turned green and showed differentiation of globular and heart embryos when transferred to the modified B5 medium, without formation of shoot. These findings showed that the applied concentration of ABA may cause inhibition of conversion of globular and heart embryos to shoot. The increased level of BA, however, was not able to ameliorate the effect of ABA. The effects of two types of auxin (IBA and NAA), an auxin pretreatment and the physical form of the culture medium on in vitro rooting of micro shoots of tea (*Camellia sinensis* L.) was studied (Gunasekare and Evans, 2000). The incorporation of auxin in the rooting medium did not bring about the formation of roots but caused basal callus formation. Microshoots were successfully induced to root with a pretreatment of IBA (50 mg/litre dip for 3 h) followed by culture of shoots in auxin-free half-strength MS liquid medium with continuous agitation. Shake cultures accelerated the and increased the number and the length of the roots compared to static cultures. An initial dark period of 7 or 14 days had no significant effect on root initiation, over cultures exposed to a continuous 12 h photoperiod. Aye *et al.* (2008) indicated the invention provides regeneration method for tea plant from leaf explants via callus phase, where in explants were cultured on basal MS medium supplemented with 3, 5, 7 mg/L 2, 4-D (2, 4-dichloro phenoxy acetic acid). Callus induction was observed by 5mg/L 2, 4-D for a period of 4-6 weeks. Shoot initiation and rhizogenesis from the leaf- derived callus of tea has been standardized using the various concentrations of 1, 3,



5, 7 mg/L BAP only and combination with 1mg/L IBA. It is now accepted that the single concentration of 3mg/L BAP was the best for shoot initiation and rhizogenesis

calli were obtained by combination of 1mg/L IBA and 3mg/L BAP.




**Figure-1.** Formation of shoot and root after in vitro culture on MS basal medium.

**Table-1.** Analysis of variance of IBA hormone levels on rooting in micro cuttings of tea.

Sours of variance	df	Root length	Root number
Replication	3	0.81	1.05
IBA hormone levels	4	13.88**	149.97**
Error	12	0.007	0.081
C.V %		3.15	3

\*\* and \* respectively significant in 1% and 5% area.



**Figure-2.** The effect of BA hormone levels on shoot length in vitro culture of Tea.



www.arnjournals.com




Figure-3. The effect of IBA hormone levels on root number in micro cuttings of tea.




Figure-4. Correlation between root number and root length in micro cuttings of tea.

## REFERENCES

Aye T.H., K.M Lwin and A.A. Khai. 2008. Propagation of Commercial Tea (*Camellia sinensis* (L.) O. Kuntze) by Efficient In Vitro Tissue Culture Method. GMSARN International Conference on Sustainable Development: Issues and Prospects for the GMS, 12-14 Nov. pp. 1-3.

Brand M.H. and R.D. Lineberger. 1986. In vitro propagation of *Halesia carolina* L. and the influence of explantation timing on initial shoot proliferation. Plant Cell, Tissue and Organ Culture. 7: 103-113.

Briggs B.A. and S.M. McCulloch. 1984. Progress in micropropagation of woody plants in the United States and western Canada. Combined Proceedings. International Plant Propagators' Society. 33: 239-248.

Debergh P.C. and P.E. Read. 1991. Micropropagation. In: Debergh PC, Zimmerman RH, (editors). The Netherlands: Kluwer Acad. Publ. pp. 1-13.

Ghanatia F. and M. Rahmati Ishkaa. 2009. Investigation of the interaction between abscisic acid (ABA) and

excess benzyladenine (BA) on the formation of shoot in tissue culture of tea (*Camellia sinensis* L.). International Journal of Plant Production. 3(4): 7-14.

Gunasekare M.T.K. and P.K. Evans. 2000. In vitro rooting of microshoots of tea (*Camellia sinensis* L.). Sri Lanka Journal of Tea Science. 66: 5-15.

Haberlandt G. 1902. Kulturversuche mit isolierten pflanzenzellen. Weisen Wien Naturwissenschaften. 111: 69-92.

2004. IAEA-TECDOC. Low cost options for tissue culture technology for developing countries. Vienna: IAEA.

Li J.R. and G.W. Eaton. 1984. Growth and rooting of grape shoot apices in vitro. Hort Science. 19: 64-65.

Lineberger R.D. 1983. Shoot proliferation, rooting, and transplant survival of tissue cultured "Hally Jolivet" cherry. Hort Science. 18: 182-185.



- Marimuthu S. and R. Raj Kumar. 2001. Physiological and Biochemical responses of micro propagated tea plants. *In Vitro Cell Dev. Biol- Plant*. 37: 618-621.
- Misawa M. 1994. Plant Tissue Culture: an alternative for production of useful metabolite. Chapter 4, 6. *FAO Agricultural Services Bulletin No. 108, Food and Agriculture Organization of the United Nations Rome, Italy*. <http://www.fao.org/docrep/t0831e/t0831e00.htm>.
- Mondal T.K. 2004. Biotechnological improvements of tea. *ISB News Report*, access entire News Report at <http://www.isb.vt.edu>.
- Murashige T. and T. Skoog. 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Plant Physiol*. 15: 473-97.
- Rout G.R., A. Mohapatra and S. Mohan Jain. 2006. Tissue culture of ornamental pot plant: A critical review on present scenario and future prospects. *Biotechnology Advances*. pp. 1-30.
- Sánchez M., S.H. Gurusinghe, K.J. Bradford and J.M. Vázquez-Ramos. 2005. Differential response of PCNA and Cdk-A proteins and associated kinase activities to benzyladenine and abscisic acid during maize seed germination. *Journal of Experimental Botany*. 56: 515-523.
- Sghaier B., W. Kriaa, M. Bahloul, J.V. Jorri'n Novo and N. Drira 2009. Effect of ABA, arginine and sucrose on protein content of date palm somatic Embryos. *Scientia Horticulturae*. 120: 379-385.
- Smart L. 2008. EFB530 Plant Physiology, Cytokinins and cell division, EFB530 Plant Physiology-Syllabus with lecture notes - spring 2008. <http://www.esf.edu/efb/course/EFB530/EFB530Syllabus.htm>.
- Steward F.C., M.O. Mapes and K. Mears. 1985. Growth and organized development of cultured cells: II. Organization in cultured grown from freely suspended cells. *Am. J. Bot.* 45: 705.
- Tahardi J.S. and W.A. Imron Riadi Dodd. 2003. Enhancement of somatic embryo development and plantlet recovery in *Camellia sinensis* by temporary liquid immersion. *Journal Bioteknologi Pertanian*. 8: 1-7.
- Tahardi J.S., I. Riyadi and W.A. Dodd. 2003. Enhancement of somatic embryo development and plantlet recovery in *Camellia sinensis* by temporary liquid immersion. *Journal Bioteknologi Pertanian*. 8(1): 1-7.
-